Contents of Vol. 27, 1990

Ahokas, H.: Absence of variation in endospermal proteins studied by electrophoresis in marginal populations of Leersia oryzoides (Poaceae), a threatened species in Finland 309–316

Apelgren, K.: Variation and distribution of Myosotis laxa sensu lato (Boraginaceae) in the Baltic region 287–299

Arts, G.H.P.: Aquatic Bryophyta as indicators of water quality in shallow pools and lakes in The Netherlands 19–32

Arup, U.: Caloplaca glomerata, a new calcicolous lichen species from Sweden 329–333

Backéus, I.: Production and depth distribution of fine roots in a boreal open bog 261–265

Botch, M.: Aapa-mires near Leningrad at the southern limit of their distribution 281–286

Castrén, M., Haavisto, T., Leinonen, K.P., Poutanen, H. & Nuorteva, P.: The influence of soil preparation on the concentrations of iron, aluminium, zinc and cadmium in leaves of Betula spp. and Maianthemum bifolium 139–143

Cela Renzoni, G., Viegi, L., Stefani, A. & Onnis, A.: Different in vitro germination responses in Pinus pinea pollen from two localities with different levels of pollution 85–90

Enroth, J.: Bryophyte flora of the Huon Peninsula, Papua New Guinea. XXXVIII. Cyrtopodaceae and Cryphaeaceae (Musci) 177–188

Grolle, R. & Piippo, S.: Bryophyte flora of the Huon Peninsula, Papua New Guinea. XXXVII. Leucolejeunea (Lejeuneaceae, Hepaticae) 119–129

Harmaja, H.: On the taxonomy and chorology of Carex pallens 79–83

Harmaja, H.: New names and nomenclatural combinations in Rhododendron (Ericaceae) 203–204

Harmaja, H. & Pellinen, K.: Three different chromosome numbers from Finnish Arabidopsis suecica (Brassicaceae) 335–336

Havas, P. & Hyvärinen, M.: Effect of cutting on the winter water economy of the Scots pine (Pinus sylvestris) 169–175

Hintikka, V.: Germination ecology and survival strategy of Rumex acetosella (Polygonaceae) on drought-exposed rock outcrops in South Finland .. 205–215

Jalas, J.: Sorbus aucuparia f. flavescentimarginata Jalas, forma nova 91

Jäppinen, J.-P. & Hotanen, J.-P.: Effect of fertilization on the abundance of bryophytes in two drained peatland forests in Eastern Finland 93–108

Kauppi, M. & Verseghy–Patay, K.: Determination of the distribution of lichen substances in the thallus by fluorescence microscopy 189–202

Koponen, A.: Tayloria rubicaulis, a new species of Splachnaceae (Musci) from the Venezuelan and Colombian Andes 43–46

Kortesharju, J., Savonen, K. & Säynätkari, T.: Element contents of raw humus, forest moss and reindeer lichens around a cement works in northern Finland 221–230

Kukkonen, I.: On the nomenclatural problems of Cyperus alternifolius 59–66

Kukkonen, I.: On the genus Eleocharis (Cyperaceae) in the Flora Iranica area, with revised infrageneric classification and nomenclature 109–117

Kukkonen, I.: The inflorescence structure of Kobresia myosuroides and related species of sect. Elyna (Cyperaceae) 159–167

Lampinen, R.: Anthyllis vulneraria subsp. danica, subsp. nova (Fabaceae), from Denmark 217–220

Lindholm, T.: Growth dynamics of the peat moss Sphagnum fuscum on hummocks on a raised bog in southern Finland 67–78

Lindholm, T. & Vasander, H.: Production of eight species of Sphagnum at Suurisuo mire, southern Finland 145–157

Norris, D.H. & Koponen, T.: Bryophyte flora of the Huon Peninsula, Papua New Guinea. XXXIII. Leskeaceae and Fabroniaceae (Musci) plus corrigenda and addenda to previous papers 1–12

Oksanen, J., Tynnyrinen, S. & Kärenlampi, L.: Testing for increased abundance of epiphytic lichens on a local pollution gradient 301–307

Piippo, S.: Bryophyte flora of the Huon Peninsula, Papua New Guinea. XXXIV. Key to the genera of Western Melanesian Hepaticae and Anthocerotae 13–18

Ravanko, O.: The taxonomic value of morphological and cytological characteristics in Oxycoccus (subgenus of Vaccinium, Ericaceae) species in Finland 235–239
Salonen, V. & Saari, V.: Generic composition of macrofungus communities on virgin mire site types in Central Finland 33–38
Simola, H., Hanski, I. & Liukkonen, M.: Stratigraphy, species richness and seasonal dynamics of plankton diatoms during 418 years in Lake Lovojärvi, South Finland 241–259
Skult, H., Hägg, M. & Djupsund, B.: Seasonal variation of isozyme and total protein phenotypes in populations of the lichen Parmelia omphalodes (Ascomycetes) 47–52
Sonck, C.E.: Taraxacum sagii (sect. Palustria), espèce nouvelle de France .. 39–41
Sonck, C.E.: Espèces nouvelles de Taraxacum de France III. Hautes Alpes. [New species of Taraxacum from France III. Hautes Alpes] 267–275
Sonck, C.E.: A new Taraxacum species, T. castellanum, from New Zealand 277–279
Stenroos, S. & Ahti, T.: The lichen family Cladoniaceae in Tierra del Fuego: problematic or otherwise noteworthy taxa 317–327
Takala, K., Olkkonen, H., Jääskeläinen, J. & Selkäinaho, K.: Total chlorine content of epiphytic and terricolous lichens and birch bark in Finland . 131–137
Tonteri, T. & Haila, Y.: Plants in a boreal city: Ecological characteristics of vegetation in Helsinki and its surroundings, southern Finland 337–352
Vilkka, L., Aula, I. & Nuorteva, P.: Comparison of the levels of some metals in roots and needles of Pinus sylvestris in urban and rural environments at two times in the growing season 53–57
Warner, B.G. & Chinnappa, C.C.: Pollen polymorphism in Empetrum nigrum (Empetraceae) in Finland 231–234

Baltic Sea Phytoplankton Identification Sheet:

4. Kuosa, H. & Hällfors, G.: Koliella spiralis Kuosa (Chlorophyceae, Ulotrichales) 353–354

Book Reviews:

Alstrup, V. & Hawksworth, D.L. 1990: The lichenicolous fungi in Greenland. (S. Huhtinen) 138
Bischler–Causse, H. 1989: Marchantia L. The Asiatic and Oceanic taxa. (S. Piippo) 130
Ehrendorfer, F. (ed.) 1989: Woody plants — evolution and distribution since the Tertiary. (T. Koponen) 84
Hämet-Ahti, L., Palmén, A., Alanko, P. & Tigerstedt, P.M.A. 1989: Suomen puu- ja pensaskasvio. (Summary: Woody flora of Finland.) (S. Hinneri) 266
Kinzel, H. 1989: Stoffwechsel der Zelle. Die zentralen Vorgänge des Stoffwechsels mit ihren physikalisch-chemischen Grundlagen. (L. Simola) 280
Øllgaard, B. 1989: Index of the Lycopodiaceae. (I. Kukkonen & J. Sarvela) 240
Renner, S.S. 1989: Systematic studies in the Melastomataceae: Bellucia, Loreya, and Macairea. (H. Tuomisto) 118
Simpson, B.B.: Krameriaceae. — Flora Neotropica. Monograph 49 (H. Tuomisto) 42
Walters, S.M., Alexander, J.C.M., Brady, A., Brickell, C.D., Cullen, J., Green, P.S., Heywood, V.H., Matthews, V.A., Robson, N.K.B., Yeo, P.F. & Knees, S.G. (eds.) 1989: The European Garden Flora. III. (L. Hämet-Ahti) 260

Numbers distributed

Vol. 26(4):349–440 ..... 17 November 1989
Vol. 27(1):1–91 ..... 7 May 1990
Vol. 27(2):93–204 ..... 14 June 1990
Vol. 27(3):205–300 ..... 3 December 1990